arXiv:1303.2222v1 [math.AG] 9 Mar 2013

THE FIRST COHOMOLOGY OF SEPARABLY RATIONALLY CONNECTED VARIETIES

FRANK GOUNELAS

Over an algebraically closed field of characteristic zero, a smooth projective separably rationally connected variety X has $H^i(X, \mathcal{O}_X) = 0$ for $i > 0$ from Hodge theory (see [Deb03] page 249). In positive characteristic we prove that at least $H^1(X, \mathcal{O}_X) = 0$, which is a special case of the following

Theorem. *Let X be a smooth projective variety over an algebraically closed field k and $f : C \to X$ a morphism from a smooth projective curve such that f^*T_X is an ample bundle. Then $H^1(X, \mathcal{O}_X) = 0$.*

Exactly as in the case of $C = \mathbb{P}^1$, one proves $H^0(X, \Omega^m_X) = 0$ for $m > 0$. Over $\mathbb{C}$ the theorem follows as before from Hodge theory. The main structure of our proof in positive characteristic follows mutatis mutandis from the proof of Theorem 2.1 in Nygaard's paper [Nyg78]. First note that we may restrict to the algebraic closure of a finite field by spreading out over a suitable finitely generated $\mathbb{F}_p$-algebra A and noting that $f : C \to X$ having f^*T_X as an ample bundle is an open condition in the setting over $\operatorname{Spec} A$. Hence assume $f : C \to X$ with f^*T_X ample is defined over $\bar{\mathbb{F}}_p$. Consider now the Artin-Schreier sequence of étale sheaves on X

$$0 \to \mathbb{F}_p \to \mathbb{G}_a \xrightarrow{F-1} \mathbb{G}_a \to 0. \tag{1}$$

The cohomologies of $\mathbb{G}_a$ and $\mathcal{O}_X$ agree and since the latter is coherent, étale and Zariski cohomology agree hence we may assume that all cohomology is in the étale site. We obtain an exact sequence

$$0 \to H^1(X, \mathbb{F}_p) \to H^1(X, \mathcal{O}_X) \xrightarrow{F-1} H^1(X, \mathcal{O}_X) \to 0 \tag{2}$$

where the last map is surjective due to SGA7.XXII Proposition 1.2. Suppressing base points, we use a method of Suwa [Suw83] to prove that a p-Sylow subgroup in the étale fundamental group $\pi_1(X)$ is trivial (in the case of $C = \mathbb{P}^1$ Kollàr has proved that $\pi_1(X)$ is trivial, using the de Jong-Starr Theorem, see [Deb03] Corollaire 3.6). Suwa, using a computation in crystalline cohomology, first proves that the vanishing of global differential forms implies that $h^i_p = \dim H^i(X, \mathbb{Q}_p) = 0$ for $i > 0$ from which $\chi_p(X) = \sum_i (-1)^i h^i_p = 1$. The proof is identical in our setup. Now pulling back $f : C \to X$ under an étale cover $Y \to X$ gives a smooth projective curve (possibly of higher genus) $g : C' \to Y$ with g^*T_Y also ample. If $\tilde{X} \to X$ the universal covering and $Y \to X$ an étale cover corresponding to a p-Sylow in $\pi_1(X)$ then $\tilde{X} \to Y$ is étale of degree p^r for some r. By Crew's formula, $\chi_p(\tilde{X}) = p^r \chi_p(Y)$ hence $p^r = \deg(\tilde{X}/Y) = 1$. Hence $\pi_1(X)$ has no p-Sylow and so $\pi_1^{ab}(X) \otimes \mathbb{F}_p = 0$.

Now $H^1(X, \mathbb{F}_p) = \operatorname{Hom}(\pi_1(X), \mathbb{F}_p) = \operatorname{Hom}(\pi_1^{ab}(X) \otimes \mathbb{F}_p, \mathbb{F}_p) = 0$ and by SGA7.XXII Proposition 2.25, the semi-simple component of $H^1(X, \mathcal{O}_X)$ under the endomorphism induced by Frobenius F is isomorphic to $H^1(X, \mathbb{F}_p) \otimes \bar{\mathbb{F}}_p$, which is trivial. Hence F is nilpotent on $H^1(X, \mathcal{O}_X)$. The injectivity of the map of the corresponding sheaves induces $H^0(X, \mathcal{O}_X/F\mathcal{O}_X) \to H^0(X, \Omega^1_X) = 0$ and so from the cohomology of the short exact sequence

$$0 \to \mathcal{O}_X \xrightarrow{F} \mathcal{O}_X \to \mathcal{O}_X/F\mathcal{O}_X \to 0 \tag{3}$$

we obtain that $F : H^1(X, \mathcal{O}_X) \to H^1(X, \mathcal{O}_X)$ is injective. Since F is thus injective and nilpotent on first cohomology, the result follows.

Acknowledgements. The author would like to thank Jakob Stix for helpful conversations.

References

[Deb03] Olivier Debarre. Variétés rationnellement connexes (d'après T. Graber, J. Harris, J. Starr et A. J. de Jong). *Astérisque*, (290):Exp. No. 905, ix, 243–266, 2003. Séminaire Bourbaki. Vol. 2001/2002.

[Nyg78] Niels Nygaard. On the fundamental group of a unirational 3-fold. *Invent. Math.*, 44(1):75–86, 1978.

[Suw83] Noriyuki Suwa. A note on the fundamental group of a unirational variety. *Proc. Japan Acad. Ser. A Math. Sci.*, 59(3):98–99, 1983.

Frank Gounelas, Mathematisches Institut, Universität Heidelberg, Im Neumheimer Feld 288, 69120, Heidelberg.

Date: March 12, 2013.